

Almond IPM: Monitoring carob moth in almonds - Update

INTRODUCTION

Pest monitoring is a critical component of IPM systems. It can provide growers with valuable information on the presence, distribution and population levels of pests, and on the impact of pest management activities. Monitoring of carob moth in almonds is documented in detail in the Australian Almonds fact sheet 'Carob moth in almonds: monitoring guidelines'. This update provides further information including findings from more recent research and is intended to help growers achieve better results from their monitoring efforts.

NUT SAMPLING

Summer sampling of new crop nuts

Sampling of the new crop of nuts in summer is used to determine infestation and damage levels to help in the planning of post-harvest management of the crop. Orchard blocks that show higher levels of pest infestation may be targeted for earlier harvest and fumigation and/or processing, to limit further damage.

Sampling position is important

Nuts collected from the ground immediately after trees are shaken represent the entire crop throughout each tree, and so provide a good overall average of damage or infestation levels throughout the tree. Nuts collected from trees before harvest represent only those areas from which they are collected and may yield infestation and damage levels that differ markedly from the overall average, as is the case with carob moth.

Carob moth is known to generally infest almonds only after hull split (Gothilf 1984; Madge et al 2015), and hull split is known to start in the upper parts of trees (Flint 2002). It could therefore be expected that higher levels of damage would be found in those upper areas where nuts are exposed to infestation for the longest period prior to harvest. Detailed sampling in the SA Riverland and Robinvale districts during the 2018, 2019 and 2020 harvests confirmed this to be the case. As Figure 1 shows, nuts taken from the upper third of the canopy of 6 m high trees had much higher damage levels compared to those from the middle and lower thirds of the canopy. The 'Lower canopy' samples represent nuts that could be picked by someone standing on the ground.

This pattern was observed in five blocks that were sampled during the study, where kernel damage levels ranged from 3 to 18% in the upper canopy and 2 to 9% overall. Across the five blocks, samples picked by hand from ground level would have underrepresented the average damage levels of the trees by between 45% and 90%. In contrast, and as expected, kernel damage levels in samples collected from the ground after tree-shake, provided much more accurate measures of average damage levels throughout the trees.




Figure 1. Variation in carob moth damage to almond kernels at harvest between various sampling positions (mean of two blocks, 2019).

Take-home message: To determine carob moth damage levels in new crops, sample nuts from the ground after tree-shake, or ensure that the full height of the canopy is represented in samples collected directly from trees.

Timing of sampling is important

Local and overseas experience has shown that carob moth infests almonds as soon as hull split occurs; kernel damage is seen within two weeks; and damage can continue to increase rapidly until the nuts are harvested and disinfested or processed. As Figure 2 shows, the percent of kernels damaged can more than double over a two-week period. Pre-harvest samples collected a considerable time (e.g. 2-3 weeks) before harvest may lead to a significant under-estimation of actual crop damage levels at harvest.




Figure 2. Growth in kernel damage by carob moth over time.

Take-home message: Pre-harvest samples to determine damage levels from carob moth should be collected as close as practicable to the time of harvest.

Winter sampling of mummy nuts

Sampling of old mummy nuts and leftover crop from the most recent harvest can be used to determine the levels of carob moth infestation being carried into winter. This may help orchard managers to target hygiene efforts towards the most heavily infested blocks.

Sampling position appears unimportant

Extensive sampling of mummy nuts from high and low in tree canopies, and from the ground, over two winters, showed that on average, levels of infestation with carob moth varied very little between those sampling positions (Figure 3).




Figure 3. Variation in infestation of almond mummy nuts with live carob moth between various sampling positions.

Take-home message: When sampling mummy nuts for carob moth infestation in winter, the position from which nuts are collected does not appear to be important.

MOTH TRAPPING

Pheromone-based male traps are useful indicators of carob moth flight patterns and population levels as described in the monitoring guidelines mentioned above.

Trap position influences the result

Like some other moth pests of tree crops, higher levels of carob moth activity tend to be observed higher in the tree canopy. Because of this, traps placed nearer the top of the canopy tend to capture more carob moths than traps placed lower in the canopy, such as at head-height (Figure 4). However, although high traps have been observed to catch up to 50% more moths than low traps, low traps still provide a useful guide to relative levels of carob moth activity over time and between orchard blocks.




Figure 4. Carob moth catches in high and low traps in almonds.

Take-home message: High-placed pheromone traps capture more carob moths, but low traps are probably just as effective for determining relative levels of moth activity.

REFERENCES

- Flint M.L. (2002). Integrated pest management for almonds, 2nd Edition. University of California, Division of Agriculture and Natural Resources. 199pp.
- Gothilf S. (1984). Biology of *Spectrobrates ceratoniae* on almonds in Israel. *Phytoparasitica* 12(2): 77-87.
- Madge D.G., Taylor C., Williams D.G. (2015). Managing carob moth in almonds. Final Report, Project AL12004. Hort Innovation, Sydney, Australia.

Prepared by Agriculture Victoria Research (Invertebrate & Weed Sciences) for Hort Innovation project AL16009 'An Integrated Pest Management program for the Australian almond industry'.

DISCLAIMER This publication may be of assistance to you but the State of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

© The State of Victoria Department of Jobs, Precincts and Regions, October 2020.



Strategic levy investment



This project has been funded by Hort Innovation using the almond research and development levy and funds from the Australian Government. For more information on the fund and strategic levy investment visit horticulture.com.au